

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/316619650>

Space-Time Evolution of Seismicity in The North-Western Geysers Geothermal Field and its Connection to Stimulatio....

Poster · April 2017

CITATIONS

0

READS

117

5 authors, including:



Konstantinos Michail Leptokaropoulos

Polish Academy of Sciences

50 PUBLICATIONS 56 CITATIONS

[SEE PROFILE](#)



Monika Staszek

Polish Academy of Sciences

15 PUBLICATIONS 28 CITATIONS

[SEE PROFILE](#)



Stanislaw Lasocki

Polish Academy of Sciences

83 PUBLICATIONS 495 CITATIONS

[SEE PROFILE](#)



Grzegorz Kwiatek

Helmholtz-Zentrum Potsdam - Deutsches Ge...

73 PUBLICATIONS 752 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Digital Research Space of Induced Seismicity for EPOS Purposes (IS-EPOS project) [View project](#)



Nova project Aspo [View project](#)


Space-Time Evolution of Seismicity in The North-Western Geysers Geothermal Field and its Connection to Stimulation Processes

Konstantinos Leptokaropoulos¹, Monika Staszek¹, Stanislaw Lasocki¹, Grzegorz Kwiatek², Patricia Martínez-Garzón²

¹Seismology and Physics of the Earth's Interior, Institute of Geophysics, Polish Academy of Sciences, Warsaw, Poland, kleptoka@igf.edu.pl.


²Helmholtz-Centre Potsdam GFZ German Research Centre for Geosciences, Potsdam, Germany

Introduction: Hydraulic fracturing that has been carried out during the last decades is occasionally connected with triggered and induced seismicity worldwide. Data from geothermal fields can be used as a proxy for shale gas exploitation associated seismicity since they are both tightly connected to hydraulic fracturing. The Geysers (California, USA) is the largest producing geothermal field in the world and has been sufficiently monitored and studied since the last decades. Seismic and technological data gathered during the past 40 years indicate a connection of seismic activity with the fluctuations of the injected fluid volume (Majer & Peterson, 2007; Martinez-Garzon and Kwiatek, 2014; Johnson et al., 2016).




Data: We utilize 'The Geysers' seismic and water injection data from the compiled SHEER (Shale Gas Exploration and Exploitation Induced Risks) database. In this study we verify and quantify this correlation by analyzing the data associated with 2 injection wells (Prati-9 and Prati-29) which covers a time period of approximately 7 years (1154 events with $M \geq 1.4$, occurred from November 2007 to August 2014), as they were relocated by Kwiatek et al., 2015.

Acknowledgements: This work was supported within SHEER: "Shale Gas Exploration and Exploitation Induced Risks" project funded from Horizon 2020 – R&I Framework Programme, call H2020-LCE-2014-1 and within statutory activities No3841/E-41/S/2016 of Ministry of Science and Higher Education of Poland.




Methods: The correlation between spatio-temporal seismicity evolution and variation of the injection data is performed by elaboration of original and smoothed time-series through specified statistical tools (cross correlation, binomial test to investigate significant rate changes, b-value variation).



References:

- Johnson C. W., Totten E. J., Bürgmann R. (2016), Depth migration of seasonally induced seismicity at The Geysers geothermal field, *Geophysical Research Letters*, 43, doi:10.1002/2016GL069546.
- Kwiatek, G., P. Martínez-Garzón, G. Dresen, M. Bohnhoff, H. Sone and C. Hartline (2015), Effects of long-term fluid injection on induced seismicity parameters and maximum magnitude in northwestern part of the Geysers geothermal field, *J. Geophys. Res. Solid Earth*, 120, doi:10.1002/2015JB012362.
- Leptokaropoulos K., Staszek M., Lasocki S., Martínez-Garzón P and Kwiatek G., Space-Time Evolution of Seismicity and its Connection to Stimulation Processes, in North-Western Geysers Geothermal Field (manuscript under preparation).
- Majer, E. L., and J. E. Peterson (2007), The Impact of Injection on Seismicity at The Geysers, California Geothermal Field., *Int. J. Rock Mech. Min.*, 44, 1079-1090.
- Martínez-Garzón, P., G. Kwiatek, H. Sone, M. Bohnhoff, G. Dresen, and C. Hartline (2014), Spatiotemporal changes, faulting regimes, and source parameters of induced seismicity: A case study from The Geysers geothermal field. *J. Geophys. Res. Solid Earth*, 119, 8378-8396.



Results: Our analysis indicate a short time delay of seismicity occurrence at distances larger than 200m from the injection well, whereas no evidence of significant correlation between injection rates and b-values was discovered. The obtained results are expected to contribute to a better comprehension of the role of hydraulic fracturing and the physical processes that control seismogenesis in fluid injection sites (Leptokaropoulos et al., manuscript under preparation).